

Taylor Series

Recall writing a function as a Power Series:

$$f(x) = c_0 + c_1(x - a) + c_2(x - a)^2 + c_3(x - a)^3 + \dots + c_n(x - a)^n + \dots$$

for $x = a$, $f(a) =$

so, $c_0 =$

$$f'(x) = c_1 + 2c_2(x - a) + 3c_3(x - a)^2 + 4c_4(x - a)^3 + \dots + nc_n(x - a)^{n-1} + \dots$$

then, $f'(a) =$

so, $c_1 =$

$$f''(x) = 2c_2 + 2 \cdot 3c_3(x - a) + 3 \cdot 4c_4(x - a)^2 + \dots + (n - 1)n \cdot c_n(x - a)^{n-2} + \dots$$

then, $f''(a) =$

so, $c_2 =$

$$f'''(x) = 2 \cdot 3c_3 + 2 \cdot 3 \cdot 4c_4(x - a) + \dots + (n - 2)(n - 1)n \cdot c_n(x - a)^{n-3} + \dots$$

then, $f'''(a) =$

so, $c_3 =$

Do you see a pattern?

Predict $f^{(n)}(a) =$

so, $c_n =$

Therefore, a function as a Power Series:

$$f(x) = c_0 + c_1(x - a) + c_2(x - a)^2 + c_3(x - a)^3 + \dots + c_n(x - a)^n + \dots$$

can now be written as the

Taylor Series

$$f(x) = \quad + \quad (x - a) + \quad (x - a)^2 + \quad (x - a)^3 + \dots + \quad (x - a)^n + \dots$$

$$= \sum_{k=0}^{\infty} \quad (x - a)^k$$

where f is generated (centered) @ $x = a$

and the partial sum, called the

Taylor Polynomial of order n for $f(x)$

$$P_n(x) = 1 + (x - a) + \frac{(x - a)^2}{2!} + \frac{(x - a)^3}{3!} + \dots + \frac{(x - a)^n}{n!}$$
$$= \sum_{k=0}^n \frac{(x - a)^k}{k!}$$

where f is generated (centered) @ $x = a$

For a more specific case, we have:

Maclaurin Series

$$f(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$
$$= \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

where f is generated (centered) @ $x = 0$

Example 1:

Find the Maclaurin Series of $f(x) = e^x$

Example 2:

Find the Maclaurin Series of $f(x) = \sin x$