

AP[®] Calculus BC 2010 Free-Response Questions

The College Board

The College Board is a not-for-profit membership association whose mission is to connect students to college success and opportunity. Founded in 1900, the College Board is composed of more than 5,700 schools, colleges, universities and other educational organizations. Each year, the College Board serves seven million students and their parents, 23,000 high schools, and 3,800 colleges through major programs and services in college readiness, college admission, guidance, assessment, financial aid and enrollment. Among its widely recognized programs are the SAT®, the PSAT/NMSQT®, the Advanced Placement Program® (AP®), SpringBoard® and ACCUPLACER®. The College Board is committed to the principles of excellence and equity, and that commitment is embodied in all of its programs, services, activities and concerns.

© 2010 The College Board. College Board, ACCUPLACER, Advanced Placement Program, AP, AP Central, SAT, SpringBoard and the acorn logo are registered trademarks of the College Board. Admitted Class Evaluation Service is a trademark owned by the College Board. PSAT/NMSQT is a registered trademark of the College Board and National Merit Scholarship Corporation. All other products and services may be trademarks of their respective owners. Permission to use copyrighted College Board materials may be requested online at: www.collegeboard.com/inquiry/cbpermit.html.

Visit the College Board on the Web: www.collegeboard.com. AP Central is the official online home for the AP Program: apcentral.collegeboard.com.



CALCULUS AB SECTION II, Part A

Time—45 minutes
Number of problems—3

A graphing calculator is required for some problems or parts of problems.

1. There is no snow on Janet's driveway when snow begins to fall at midnight. From midnight to 9 A.M., snow accumulates on the driveway at a rate modeled by $f(t) = 7te^{\cos t}$ cubic feet per hour, where t is measured in hours since midnight. Janet starts removing snow at 6 A.M. (t = 6). The rate g(t), in cubic feet per hour, at which Janet removes snow from the driveway at time t hours after midnight is modeled by

$$g(t) = \begin{cases} 0 & \text{for } 0 \le t < 6\\ 125 & \text{for } 6 \le t < 7\\ 108 & \text{for } 7 \le t \le 9 \end{cases}.$$

(a) How many cubic feet of snow have accumulated on the driveway by 6 A.M.?

(b) Find the rate of change of the volume of snow on the driveway at 8 A.M.

Do not write beyond this border.

Do not write beyond this border

(c) Let h(t) represent the total amount of snow, in cubic feet, that Janet has removed from the driveway at time t hours after midnight. Express h as a piecewise-defined function with domain $0 \le t \le 9$.

(d) How many cubic feet of snow are on the driveway at 9 A.M.?

Do not write beyond this border.

GO ON TO THE NEXT PAGE.

Do not write beyond this border



2



2









t (hours)	0	2	5	7	8
E(t) (hundreds of entries)	0	4	13	21	23

- 2. A zoo sponsored a one-day contest to name a new baby elephant. Zoo visitors deposited entries in a special box between noon (t = 0) and 8 P.M. (t = 8). The number of entries in the box t hours after noon is modeled by a differentiable function E for $0 \le t \le 8$. Values of E(t), in hundreds of entries, at various times t are shown in the table above.
 - (a) Use the data in the table to approximate the rate, in hundreds of entries per hour, at which entries were being deposited at time t = 6. Show the computations that lead to your answer.

(b) Use a trapezoidal sum with the four subintervals given by the table to approximate the value of $\frac{1}{8} \int_0^8 E(t) dt$. Using correct units, explain the meaning of $\frac{1}{8} \int_0^8 E(t) dt$ in terms of the number of entries.

(c) At 8 P.M., volunteers began to process the entries. They processed the entries at a rate modeled by the function P, where $P(t) = t^3 - 30t^2 + 298t - 976$ hundreds of entries per hour for $8 \le t \le 12$. According to the model, how many entries had not yet been processed by midnight (t = 12)?

(d) According to the model from part (c), at what time were the entries being processed most quickly? Justify your answer.

Do not write beyond this border.

The beyond this porder.



- 3. A particle is moving along a curve so that its position at time t is (x(t), y(t)), where $x(t) = t^2 4t + 8$ and y(t) is not explicitly given. Both x and y are measured in meters, and t is measured in seconds. It is known that $\frac{dy}{dt} = te^{t-3} 1$.
 - (a) Find the speed of the particle at time t = 3 seconds.

(b) Find the total distance traveled by the particle for $0 \le t \le 4$ seconds.

Do not write beyond this border.

Continue problem 3 on page 9.

Do not write beyond this border.



Do not write beyond this border

(c) Find the time t, $0 \le t \le 4$, when the line tangent to the path of the particle is horizontal. Is the direction of motion of the particle toward the left or toward the right at that time? Give a reason for your answer.

- (d) There is a point with x-coordinate 5 through which the particle passes twice. Find each of the following.
 - (i) The two values of t when that occurs
 - (ii) The slopes of the lines tangent to the particle's path at that point
 - (iii) The y-coordinate of that point, given $y(2) = 3 + \frac{1}{e}$

END OF PART A OF SECTION II

IF YOU FINISH BEFORE TIME IS CALLED, YOU MAY CHECK YOUR WORK ON
PART A ONLY. DO NOT GO ON TO PART B UNTIL YOU ARE TOLD TO DO SO.

NO CALCULATOR ALLOWED


CALCULUS AB

SECTION II, Part B


Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.



- 4. Let R be the region in the first quadrant bounded by the graph of $y = 2\sqrt{x}$, the horizontal line y = 6, and the y-axis, as shown in the figure above.
 - (a) Find the area of R.



Continue problem 4 on page 11.

(b) Write, but do not evaluate, an integral expression that gives the volume of the solid generated when R is rotated about the horizontal line y = 7.

(c) Region R is the base of a solid. For each y, where $0 \le y \le 6$, the cross section of the solid taken perpendicular to the y-axis is a rectangle whose height is 3 times the length of its base in region R. Write,

Do not write beyond this border.

but do not evaluate, an integral expression that gives the volume of the solid.

NO CALCULATOR ALLOWED

- 5. Consider the differential equation $\frac{dy}{dx} = 1 y$. Let y = f(x) be the particular solution to this differential equation with the initial condition f(1) = 0. For this particular solution, f(x) < 1 for all values of x.
 - (a) Use Euler's method, starting at x = 1 with two steps of equal size, to approximate f(0). Show the work that leads to your answer.

(b) Find $\lim_{x\to 1} \frac{f(x)}{x^3-1}$. Show the work that leads to your answer.

Do not write beyond this border.

Continue problem 5 on page 13.

Do not write beyond this border

NO CALCULATOR ALLOWED

(c) Find the particular solution y = f(x) to the differential equation $\frac{dy}{dx} = 1 - y$ with the initial condition f(1) = 0.

Do not write beyond this border.

盏..

Do not write beyond this borde



.6



6











$$f(x) = \begin{cases} \frac{\cos x - 1}{x^2} & \text{for } x \neq 0 \\ -\frac{1}{2} & \text{for } x = 0 \end{cases}$$

6. The function f, defined above, has derivatives of all orders. Let g be the function defined by

$$g(x) = 1 + \int_0^x f(t) dt.$$

(a) Write the first three nonzero terms and the general term of the Taylor series for $\cos x$ about x = 0. Use this series to write the first three nonzero terms and the general term of the Taylor series for f about x = 0.

(b) Use the Taylor series for f about x = 0 found in part (a) to determine whether f has a relative maximum, relative minimum, or neither at x = 0. Give a reason for your answer.

(c) Write the fifth-degree Taylor polynomial for g about x = 0.

(d) The Taylor series for g about x = 0, evaluated at x = 1, is an alternating series with individual terms that decrease in absolute value to 0. Use the third-degree Taylor polynomial for g about x = 0 to estimate the value of g(1). Explain why this estimate differs from the actual value of g(1) by less than $\frac{1}{6!}$.

To not with obyond this bolder